

Who to contact if you are worried or require further information

For general enquiries about appointments, please phone:

Mr Smibert's secretary on: 01935 384597

Mr Chambler's secretary on: 01935 384779

For enquiries regarding admission dates, please phone the Admissions Office on: 01935 384619

If your wound changes in appearance, weeps fluid or pus, or you feel unwell with a high temperature, contact your GP.

If you have a query about **exercises or movements**, contact the **Physiotherapy department** where you are having treatment if you have already started, or else the Yeovil Physiotherapy Department on 01935 384358.

For queries regarding **self care** (eg, dressing, bathing) contact the **Occupational Therapy Department** on 01935 384215.


Patients may be contacted by post and requested to complete and return a questionnaire. This is useful for the surgical team, for monitoring patients progress and may save an unnecessary visit to the hospital.

We would like to thank the Nuffield Orthopaedic Centre (Upper Limb Clinic) for allowing us to re-produce the information in this leaflet.

If you require this leaflet in any other format, eg, large print, please telephone 01935 384590

Information for you after your operation

Rotator Cuff Repair



This information booklet has been produced to help you gain the maximum benefit after your operation. **It is not a substitute** for professional medical care and should be used in association with treatment at the Orthopaedic Clinic. **Individual variations** requiring specific instructions not mentioned here may be required.

This booklet was compiled by:
Jane Moser (Senior Physiotherapist)
Professor Andrew Carr (Orthopaedic Surgeon)
Louise Le Good (Senior Occupational Therapist)
At the Nuffield Orthopaedic Centre, Oxford.

Help and feedback was given from people who have had rotator cuff repair surgery.

Standing facing a wall. Walk hand up the wall, try and keep shoulder down. Then take hand away from the wall and try and keep it there. Hold this for 5 seconds. Try this with hand at different heights.

Repeat 5 times (each position).



Standing with arms behind your back. Grasp the wrist of your operated arm. Gently slide your hands up your back.

Repeat 5 times. Do not force.



Stand or sit. Try and set up a pulley system with the pulley or ring high above you. Pull down with your **unoperated** arm to help lift the operated arm up.

Repeat 10 times.

These exercises are a few examples, others may be selected instead of, or in addition to these.




Phase 3 Exercises

These will concentrate on increasing the strength and mobility around your shoulder. The exercises will be selected for your individual shoulder and your lifestyle. Work hard at them as improvements in strength can increase for up to 2 years!

Phase 2 Exercise


Only start these when advised to do so.
This may not be for 6 weeks.

Lying on your back, elbow bent.
Help the operated arm up as
before but once the arm is
vertical, try and keep it there
without support of the other arm.
Gradually lower and raise the arm
in an arc, until you can lift it from
the bed.




Repeat 5-10 times. (Shown for left shoulder).

After two weeks you may be able
to progress this exercise, by doing
the same action standing up. Build
it up gradually and try and get your
shoulder joint to move.



Lying on your back, elbow bent and
close to side. Hold wrist of your
affected arm with your good hand.
Try and move your hand on the
operated arm inwards but **prevent
it from moving** by using the other hand.
Hold for 5 seconds.



Repeat 20 times. (This exercise can
also be done standing or sitting.)

In the same position, try and move
hand outwards without allowing
movement. Hold for 5 seconds.



Repeat 20 times. (This exercise can
also be done standing or sitting.)

Contents

| | Page |
|--|-------------|
| The rotator cuff | 4 |
| About the rotator cuff operation | 5 |
| The risks & complications | 5 |
| Common questions about ... | |
| a) pain | 6 |
| b) the sling | 6 |
| c) exercises | 7 |
| d) wound care | 7 |
| e) returning to the hospital | 7 |
| f) things to avoid | 8 |
| g) how you may progress | 9 |
| h) return to work | 10 |
| i) return to driving | 10 |
| j) leisure activities | 10 |
| Guide to daily activities | 11 |
| Exercises | 12 |
| Contact points for further information | 16 |



The 'Rotator Cuff'

The rotator cuff is a group of muscles closely wrapped around the shoulder. These muscles help keep the joint in the correct position and control shoulder movements. They attach from the shoulder blade onto the top of the arm bone. The muscles can be torn through general wear and tear or after an accident/fall.

The damage usually occurs close to where the muscle joins the bone (called the 'tendon'). If one or more of these muscles is torn, movement is no longer smoothly controlled and the shoulder becomes weak and painful. See pictures below.

Left shoulder, viewed from the back

Please note: in your body there are more muscles on top of these muscles and bones!



Elbow exercise


Standing or lying
Straighten your elbow and then bend your elbow.

Repeat 5 times. (Shown for left arm)



Shoulder blade exercise

Sit or stand.
Shrug shoulders up and forwards.
Then roll them down and back.
Repeat 10 times.



Shoulder exercises

These are important to do


Stand, lean forwards.
Let your arm hang down. Swing arm forward and backwards.
Keep it RELAXED.

Repeat 10 times. (Shown for left arm.)




Lying on your back on bed or floor.
(Shown for left arm.)

Lift your operated arm with your other arm. Keep the operated arm as RELAXED as possible. Can start with elbow bent.



Repeat 5 times.

Lying on back (on bed or floor),
towel under arm. Keep elbow into your side and bent. Hold stick in your hands.
Move the stick sideways, gently pushing hand on your operated arm outwards.



Repeat 5 times.

Exercises

Use pain relief medication and/or ice packs to reduce the pain before you exercise.

It is normal for you to feel aching, discomfort or stretching sensations when doing these exercises. However, if you experience intense and lasting pain (eg, more than 30 minutes), reduce the exercises by doing them less forcefully, or less often. If this does not help, discuss the problem with the physiotherapist.

Certain exercises may be changed or added for your particular shoulder.

Do short, frequent sessions (eg, 5-10 minutes, 4 times a day) rather than one long session.

Gradually increase the number of repetitions you do. Aim for the repetitions your therapist advises, the numbers stated here are rough guidelines.

Get into the habit of doing them! Good luck.

Note that all pictures are shown for the right shoulder unless specified.

Phase 1 Exercises (for 3 or 6 weeks)


Do ALL the exercises in this section unless the therapist specifically advises you not to. If you have a special arm splint, you may not be expected to do some of these exercises. You will be advised by your physiotherapist about this.

Neck exercises

Standing or sitting.

Turn your head to one side. Repeat 5 times. Then turn your head to the other side and repeat 5 times.

Tilt your head towards one shoulder. Repeat 5 times. Then tilt your head to the other side and repeat 5 times.



About your Rotator Cuff Operation

The operation is to try to re-attach the tendon(s) to the bone.

Sometimes the tear is too big and/or the tendon is too fragile for this to be possible and only a partial repair can be achieved. The repair involves sewing the torn tendon into a groove on the bone.

In addition, a ligament is also released, and a prominence on a bone cut away to give the repaired muscle more space in which to move.

Anaesthetic

The operation is performed under a general anaesthetic. Usually the anaesthetist will administer a block of the shoulder and arm beforehand, by injecting local anaesthetic into the neck. This reduces pain after surgery and makes the whole operation smoother. The benefits of this method outweigh the small risk of complications associated with this technique.

As the strength and size of the repair can vary, we can only give you rough guidelines in this booklet. The surgeon and therapists will be able to discuss your individual surgery after the operation, please ask.

What are the risks?

All operations involve an element of risk. We do not wish to over-emphasise them but feel that you should be aware of them before and after your operation. The risks include:

- a) complications relating to the **anaesthetic**
- b) **infection**. These are usually superficial wound problems. Occasionally deep infection may occur after the operation, <1%
- c) persistent **pain** and/or **stiffness** in (and around) the shoulder in 10-20% of people
- d) sometimes it is **impossible to repair the muscle** because it is too badly torn. The sub-acromial decompression should improve pain. But movement and strength may be no better.
- e) damage to **nerves** and **blood vessels** around the shoulder
- f) a need to **redo the surgery**. Sometimes the muscle can re-tear. This depends on the size of the tear and how well it heals.

Please discuss these issues with the doctors if you would like further information.

Questions that we are often asked

Will it be painful?

Although the operation is to relieve pain, it may be several weeks until you begin to feel the benefit. You will be given pain relief medication (either as tablets or injections) to help reduce the discomfort whilst you are in hospital. A prescription for continued pain relief medication will be given to you for your discharge home. If you require further medication after these are finished, please contact your general practitioner (GP).

You may find ice packs over the area helpful. Use a packet of frozen peas, placing a piece of paper towel between your skin and the ice pack. Until it is healed, also use a plastic bag to protect the wound from getting wet. Leave on for 10 to 15 minutes and you can repeat this several times a day.

Do I need to wear a sling?

Your arm will be immobilised in a sling. This is to protect the repair during the early phases of healing and to make your arm more comfortable. You will be shown how to get your arm in and out of the sling by a nurse or physiotherapist. (A further information sheet on this is available if required.)

You will wear the sling for a **minimum** of 3 weeks. If you have had a major repair you will wear the sling for 6 weeks. The sling will then gradually be used less as the repair heals and the muscles regain their strength.

You may find your armpit becomes uncomfortable whilst you are wearing the sling for long periods of time. Try using a dry pad or cloth to absorb the moisture.

If you are lying on your back to sleep, you may find placing a small towel or pillow under your upper arm will be comfortable.

Guide to activities in the first 6-12 weeks

Some difficulties are quite common, particularly in the early stages when you are wearing the sling and when you first start to take the sling off. If necessary an occupational therapist (OT) can help you and specialist equipment can be borrowed from the OT department. Below are listed some common difficulties with guides which may help.

Please discuss anything you are unsure about with the staff.

- 1. Getting on and off seats.** Raising the height can help, eg, extra cushion, raised toilet seat, chair or bed blocks.
- 2. Getting in and out of the bath.** Using bath boards can help, though initially you may prefer to strip wash.
- 3. Hair care and washing yourself.** Long handled combs, brushes and sponges can help.
- 4. Dressing.** Wear loose clothing, either with front fastening or that you can slip over your head. For ease also remember to dress your operated arm first and undress your operated arm last. In addition, dressing sticks, long handled shoe horns, elastic shoe laces and a 'helping hand' may help.
- 5. Eating.** A non-slip mat and other simple aids can help when one handed. Use your operated arm once it is out of the sling as you feel able.
- 6. Household tasks/cooking.** Do not use your operated arm for activities involving weight (eg, lifting kettle, iron, saucepan) for 8-12 weeks. Light tasks can be started once your arm is out of the sling. To begin with you may find it more comfortable keeping your elbow into your side.

You are likely to see the most progress in the first 6 months. At times it can feel like a lot of hard work for little in return. Set small, achievable goals and try and keep a positive attitude.

You may continue to see improvement in the use of your arm and shoulder for 1 to 2 years following the operation.

When can I return to work?

You may be off work between 4-12 weeks, depending on the type of job you have. If you are involved in lifting, overhead activities or manual work you are advised **not** to do these for 4 to 6 **months**. Please discuss any queries with the physiotherapist or hospital doctor.

When can I drive?

This is likely to be 2-3 weeks after you have stopped wearing the sling (ie, between 5 to 12 weeks after your operation). Check you can manage all the controls and it is advisable to start with short journeys. The seat-belt may be uncomfortable initially, but your shoulder will not be harmed by it.

In addition, check your insurance policy. You may need to inform the insurance company of your operation.

When can I participate in leisure activities?

Your ability to start these will be dependent on the pain, range of movement and strength that you have in your shoulder following the operation. Please discuss activities you may be interested in with your physiotherapist or consultant. Start with short sessions, involving little effort and gradually increase. General examples:

Gentle swimming (this will depend on the extent of the surgery)

Gardening (light tasks, eg, weeding) - after 12 weeks

Bowls, golf – after 16 weeks (4 months)

Do I need to do exercises?

Yes! To begin with you will be moving the joint **only** for specific exercises which the physiotherapist will show you. You will continue with these exercises at home for between 3-6 weeks depending on your particular operation. Out patient physiotherapy appointments will be organised to start after this. However, if your shoulder is particularly stiff or you are having problems with the exercises, you will be seen earlier.

You will need to get into the habit of doing regular daily exercises at home for several months. They will enable you to gain maximum benefit from your operation.

Some of the early exercises are shown at the back of this booklet.

What do I do about the wound and the stitches?

Keep the wound dry until it is healed. This is normally for 10 to 14 days. You can shower/wash and use ice packs but protect the wound with cling film or a plastic bag.

Avoid using spray deodorants, talcum powder or perfumes near or on the scar.

Normally your stitches or clips will be removed by the nurse at your GP surgery after 10 days. You will need to make an appointment to have this done.

When do I return to the Orthopaedic Clinic at Yeovil?

This is usually arranged for approximately 6 weeks after your discharge from hospital to check how you are progressing. Please discuss any queries or worries you have at this time.

Appointments are made after this if necessary.

Are there things that I should avoid doing?

For at least 3 weeks (possibly 6 weeks, depending on your operation)

Do not try and use the arm for everyday activities, especially those taking your elbow away from your body. Keep it in the sling, except when you are doing your exercises. Continue with this until you are told otherwise by the consultant, hospital doctor or physiotherapist.

Do not let your arm stretch across the front of your body. This can happen at night when you are lying on your unoperated side. So, once you stop using the sling, place your arm on pillows in front of you.

For 6 weeks

Do not lie on your operated side. After this time be guided by pain.

Avoid lifting any weight for 8-12 weeks (eg, a kettle). This is to avoid stressing the repaired muscle. Heavier lifting (eg, digging the garden, manual work) should be avoided for 4-6 months.

There may be other movements that are restricted for you. You will be told if this is the case.

Within these general instructions be guided by pain. It is normal for you to feel discomfort, aching and stretching sensations when you start to use your arm. Intense and aching pain (eg, for 30 minutes) is an indication to reduce that particular activity or exercise. In addition, avoid sudden, forceful movements involving weight.

How am I likely to progress?

This can be divided into three phases

Phase 1 Sling on, no movement except for exercises

You will basically be one-handed, immediately after the operation and for the first 3 weeks, or up to 6 weeks (depending on your operation). This will affect your ability to do everyday activities, especially if your dominant hand (right if you are right-handed) is the side with the operation. Activities that are affected include dressing, bathing, hair care, shopping, eating and preparing meals. Some guides to common problems are given later in this booklet (page 11). If you are having particular problems, an occupational therapist can suggest ways to help you. Before you are discharged from hospital, the staff will help you plan for how you will manage when you leave. In addition, we may be able to organise or suggest ways of getting help once you are discharged from hospital.

Phase 2 Regaining everyday movements

This starts once you are given the go-ahead by the hospital doctors or physiotherapist (between 3-6 weeks after your operation). You will have out patient physiotherapy and start exercises to gain muscle control and movement. The arm can now be used for daily activities; initially these will be possible at waist level but gradually you can return to light tasks with your arm away from your body.

Phase 3 Regaining strength

After 12 weeks you will be able to increase your activities, using your arm away from your body and for heavier tasks. The exercises now have an emphasis on regaining strength and getting maximum from your shoulder. There are still some restrictions on lifting.