

Who to contact if you are worried or require further information

For general enquiries about appointments, please phone:

Mr Smibert's secretary on: 01935 384597

Mr Chambler's secretary on: 01935 384779

For enquiries regarding admission dates, please phone the Admissions Office on: 01935 384619

If your wound changes in appearance, weeps fluid or pus, or you feel unwell with a high temperature, contact your GP.

If you have a query about **exercises or movements**, contact the **Physiotherapy department** where you are having treatment if you have already started, or else the Yeovil Physiotherapy Department on 01935 384358.

For queries regarding **self care** (eg, dressing, bathing) contact the **Occupational Therapy Department** on 01935 384215.


Patients may be contacted by post and requested to complete and return a questionnaire. This is useful for the surgical team, for monitoring patients progress and may save an unnecessary visit to the hospital.

We would like to thank the Nuffield Orthopaedic Centre (Upper Limb Clinic) for allowing us to re-produce the information

If you require this leaflet in any other format, eg, large print, please telephone 01935 384590

Information for you after your operation


Shoulder Replacement



This information booklet has been produced to help you gain the maximum benefit after your operation. **It is not a substitute** for professional medical care and should be used in association with treatment at the Orthopaedic Clinic. **Individual variations** requiring specific instructions not mentioned here may be required.

Lying on back. Try and take both hands behind neck with your elbows pointing up to the ceiling.

Then progress to allow elbows to **gently** move apart.



Phase 3 continued and Phase 4

There is great variation in what people can achieve during rehabilitation, therefore it is not possible to give all potential exercises. The physiotherapist will design an ongoing exercise programme for you which is specific to your shoulder and your needs.

Keep the exercises going until you feel there is no more improvement. This may continue for a year to 18 months – so think positive, keep at it and enjoy them!

This booklet was compiled by:
Jane Moser (Senior Physiotherapist)
Louise Le Good (Senior Occupational Therapist)
Professor Andrew Carr (Orthopaedic Surgeon)
At the Nuffield Orthopaedic Centre, Oxford.

Help and feedback was given from people who have had shoulder replacement surgery.

Start this exercise 4 weeks after your operation

Stand with arm close to side and elbow bent. Push the palm of your hand into other hand (or do this against wall, or in doorframe). Hold 10 seconds.

Aim to repeat 30 times, start with 10.




PHASE 3 EXERCISES

From 6 weeks to 12 weeks after your operation


Stand with arm close to body and elbow bent. Hold rubber exercise band. Pull your hand towards your stomach. Keep the elbow in. Control the movement on return.

Aim to repeat 30 times, start with 10.



Stand with hands clasped in front of you. Lift arms up in air. Separate your arms sideways, bend your elbows and lower them down. Progress this by:

- i) reversing the movement (start by taking elbows out to the side)



Contents

Page

The shoulder	4
Why the joint needs replacement	4
About the shoulder replacement	4-5
Risks	5
Common questions about ...	
a) pain	6
b) the sling	6
c) exercises	7
d) wound care	7
e) returning to hospital	7
f) things to avoid	8
g) how you may progress	
immediately – discharge	9
discharge – 6 weeks	9
6 weeks – 12 weeks	10
after 12 weeks	10
h) return to work	10
i) return to driving	11
j) leisure activities	12
Guide to daily activities	13
Exercises (general advice)	14
for 10 to 14 days	14-15
from 14 days to 6 weeks	16-16
from 6 weeks to 12 weeks	16-19
after 12 weeks	19
Contact points for further information	20

The Shoulder


The shoulder joint is a ball and socket joint. Most shoulder movement occurs where the ball at the top of your arm bone ('humerus') fits into the socket ('glenoid') which is part of the shoulder blade ('scapula'). See diagram below.

Why the joint needs replacement

The most common reasons for replacing the shoulder joint are for arthritis, either osteo-arthritis (wear and tear) or rheumatoid arthritis. It may also be necessary following a fracture or bad accident. With arthritis the joint becomes painful and difficult to move. Sometimes the deep layer of muscles (the 'rotator cuff') which control shoulder movements *can* also be worn or damaged.

About the shoulder replacement

The operation replaces the damaged joint surfaces. It consists of a metal replacement for the ball component and a plastic cup for the socket. See picture below of the right shoulder, viewed from the front.



Standing with elbow bent.

Help operated arm up in air with other arm. Then try and use your muscles to help, so that gradually you are able to lift the arm up without using the other arm. Try not to let the shoulder shrug up very much when you do this. You are aiming to get the shoulder joint to move smoothly.

Repeat 10 times.



Standing facing a wall. Walk hand up the wall, try and keep shoulder down. Then take hand away from wall and try and keep it there. Hold this for 5 seconds. Try this with hand at different heights.


Repeat 5-10 times.




Start this exercise 3 weeks after operation

Sitting or standing with your elbow bent. Keep elbow INTO your side. Start with your hand in front of your stomach and pull hand outwards until it is in front of you. **Control** the movement on return and do NOT try and pull out too far.


Aim to repeat 30 times, start with 10.



Standing with your operated arm against a wall. Bend your elbow. Push your HAND into the wall. Hold for 10 seconds. Repeat 30 times.




Stand with back against wall. Keep arm close to side, elbow bent. Push the elbow back into the wall. Hold for 10 seconds. Repeat 30 times.



Stand sideways with operated arm against wall. Keep arm close to side, elbow bent. Push elbow into wall. Hold for 10 seconds. Repeat 30 times.



Stand facing a wall. Keep arm close to side and elbow bent to 90°. Push your fist into the wall (use a towel if this is uncomfortable for your hand). Hold for 10 seconds.



The main reason for doing the operation is to reduce the pain in your shoulder. Ultimately, you **may** also have more movement in your shoulder. This depends on how stiff the joint was before the operation and if the muscles around the shoulder are damaged and unable to work normally. If the muscles are badly damaged, then only the ball part of the joint is replaced. When you have the operation the doctors will be able to see if there is a significant damage to the muscles and give you a realistic idea of what movements to expect.

What are the risks?

All operations involve an element of risk. We do not wish to over emphasise the risks, but feel that you should be aware of them. They include:

- a) complications relating to the **anaesthetic**
- b) **infection**. These are usually superficial wound problems. Occasionally deep infection may occur after the operation.
- c) unwanted **stiffness** and/or **pain** in (and around) the shoulder.
- d) damage to the **nerves** and **blood vessels** around the shoulder
- e) a need to **redo the surgery**

Note: Dislocation is very rare. As with all joint replacements, the components can loosen. This is not normally a problem until several years after the operation.

Please discuss these issues with the doctors if you would like further information.

Anaesthetic

The operation is performed under a general anaesthetic. Usually the anaesthetist will administer a block of the shoulder and arm beforehand, by injecting local anaesthetic into the neck. This reduces pain after surgery and makes the whole operation smoother. The benefits of this method outweigh the small risk of complications associated with this technique.

Questions that we are often asked

Will it painful?

Although the operation is to relieve pain, it may be several weeks until you begin to feel the benefit. You will be given pain relief medication (either as tablets or injections) to help reduce the discomfort whilst you are in hospital. A prescription for continued pain medication will be given to you for your discharge home. Please contact your general practitioner (GP) if you require further medication after that.

You will probably have some bruising around the shoulder/upper arm and the arm may well be swollen. This will gradually disappear over a period of a few weeks.

You may find ice packs over the area helpful. Use a packet of frozen peas, placing a piece of wet paper towel between your skin and the ice pack. Until it is healed, also use a plastic bag to protect the wound from getting wet. Leave on for 10-15 minutes and you can repeat this several times a day.

Do I need to wear a sling?

The sling is for comfort and to protect the shoulder after the operation. You can take it on and off as you wish and you do not need to have your arm strapped to your body.

The therapists and nurses will show you how to take the sling on and off. A further information sheet on how to put the sling on and off is available.

You will gradually wear the sling less over 4-8 weeks.

You may find it helpful to wear the sling at night (with or without the body strap), particularly if you tend to lie on your side. Alternatively, you can use pillows in front of you to rest your arm on.

If you are lying on your back to sleep, you may find placing a thin pillow or folded towel under your upper arm will be comfortable.


Phase 2 Exercises

From 10-14 days to 6 weeks after your operation

Lying on your back, elbow bent. Help your operated arm up as before, but once it is vertical try and keep it there without the support of the other arm.

Gradually lower and raise your arm in an arc, until you can lift it from the bed. Once this is easy, progress to standing with elbow bent (page 17).

Repeat 10 times.




Standing with arms behind your back. Grasp the wrist of your operated arm and gently stretch hand towards opposite buttock. Then slide your hands up your back.

Repeat 5 times.




Standing or sitting. Take hand of your operated arm across body towards opposite shoulder. Give gentle assistance from your other arm.


Repeat 5 times.
(Shown for left shoulder.)




Sit or stand. Shrug shoulders up and forwards.
Then roll them down and back.
Repeat 10 times.



Lying on back, folded towel under arm. **Keep elbow to side.**
Hold stick in hands. Move stick sideways, gently pushing hand on your operated arm outwards.
Do not over-stretch.
Repeat 5 times
Do not do this exercise if you have a fracture.



Lying on your back. Support your operated arm and lift up overhead.
Start with your elbows bent, then with arms straight.
Repeat 10 times. (Shown for left shoulder.)



Sit or stand. Try and set up a pulley system with the pulley or ring high above you. Pull down with your **unoperated** arm to help lift the operated arm up.
Repeat 10 times.



Do I need to do exercises?

Yes! You will be shown exercises by the physiotherapist. You will start exercises to move the shoulder on the first day after the operation. You will then need to continue with exercises when you go home and out patient physiotherapy appointments will be organised for you.

You will need to get into the habit of doing regular daily exercises at home for several months. They will enable you to gain maximum benefit from your operation.

The exercises aim to stop your shoulder getting stiff and strengthen muscles. They will be changed as you progress and made specific to your shoulder and your lifestyle. Some early exercises are shown at the back of this booklet.

What do I do about the wound?

Keep the wound dry until it is healed. This is normally for 10-14 days. You can shower or wash and use ice packs but protect the wound with cling film or plastic bag. Avoid using deodorant, talcum powder or perfumes near or on the scar.

Normally your stitches or clips will be removed by the nurse at your GP surgery after 10 days. You will need to make an appointment at the surgery to have this done.

When do I return to the Orthopaedic Clinic at Yeovil?

This is usually arranged for approximately 6 weeks after you are discharged from hospital, to check on your progress. Please discuss any queries or worries you may have when you are at the clinic. Appointments are made after this if necessary.

Patients may be contacted by post and requested to complete and return a questionnaire. This is useful for the surgical team, for monitoring patients progress and may save an unnecessary visit to the hospital.

Are there things that I should avoid?

For the first 6 weeks

1. Avoid taking your arm out to the side and twisting it backwards. For example; when putting on a shirt or coat, put your operated arm in its sleeve first. Try not to reach up and behind you (eg, seat belt in car). It is normally too painful/difficult to do! Do not force these movements for 3 months.
2. Avoid leaning with **all your body weight** on your arm with your **hand behind you**. For example, leaning heavily on your arm to get out of a chair.



The occupational therapist will show you ways of avoiding these movements and can give you aids and appliances if necessary. See 'guide to daily activities' on page 12 of this booklet.

Generally, do not be frightened to start moving the arm as much as you can. Gradually the movements will become less painful.

Exercises

General points

Use pain relief medication and/or ice packs to reduce the pain before you exercise.

It is normal for you to feel aching, discomfort or stretching sensations when doing these exercises. However, if you experience intense and lasting pain (eg more than 30 minutes), it is an indication to change the exercise by doing it less forcefully, or less often. If this does not help, discuss the problem with the physiotherapist.

Certain exercises may be changed or added for your particular shoulder.

Do short, frequent sessions (eg, 5-10 minutes, 4 times a day) rather than one long session.

Gradually increase the number of repetitions that you do. Aim for the repetitions your therapist advises, the numbers stated here are rough guidelines. After 3-4 weeks you can increase the length of time exercising.

Get into the habit of doing them! Good luck.

Note: Not all patients will be able to, or should be doing, these exercises, eg, following fractures. Your therapist will advise you.

Exercises shown for right shoulder unless specified.


Phase 1 exercises

From operation day to 10 -14 days after

Lean forwards. Let your arm hang freely. Start with small movements. Swing your arm:

- i) forwards and backwards
- ii) side to side
- iii) in circles

Repeat 5 times each movement.
(Shown for left shoulder)



Guide to daily activities in the first 4 - 6 weeks

Some difficulties are quite common, particularly in the early stages. The occupational therapist (OT) will help you to be as independent as possible during your rehabilitation. Special equipment can be borrowed from the OT department. Everyone is different so your individual needs will be assessed. We appreciate that you may have been having many of these problems before your operation. Please discuss your difficulties with the occupational therapist.

Getting on and off seats. Raising the height can help, eg, extra cushions, raised toilet seat, chair or bed blocks.

Getting in and out of the bath. Using bath boards may help. Initially you may prefer to strip wash.

Hair care and washing yourself. Long handled combs, brushes and sponges can help to stop you twisting your arm out to the side.

Dressing. Wearing loose clothing, either with front fastening or which you can slip over your head. For ease also remember to dress your operated arm first and undress your operated arm last. In addition, dressing sticks, long handled shoe horns, elastic shoe laces, sock aids and a 'helping hand' can help.

Eating. Use your operated arm as soon as you feel able for cutting up food and holding a cup. Non slip mats and other simple aids can help.

Household tasks/cooking. Do light tasks as soon as you feel able, eg, lift kettle with small amount of water, light dusting, ironing, rolling pastry. Various gadgets can help you with other tasks.

How am I likely to progress?

This can be divided into four phases

Phase 1. Immediately after the operation until you are discharged

You will start to move the shoulder with the help of the physiotherapist, but to begin with you will be quite one-handed. If your dominant hand (right hand if you are right-handed) is the side with the operation your daily activities will be affected and you will need some help.

Activities that are affected include dressing, bathing, hair care, shopping and preparing meals. The occupational therapist will discuss ways and show you how to be as independent as possible during this time. Some common difficulties which are encountered with examples of practical solutions to help are listed in a section later in this booklet (see page 12).

Before you are discharged from hospital, the staff will help you plan for how you will manage when you leave. Please discuss any worries with them. We may be able to organise or suggest ways of getting help for when you are home.

Phase 2. After you have been discharged and for up to 6 weeks after the operation

The pain in your shoulder will gradually begin to reduce and you will become more confident. Wean yourself out of the sling slowly over this time, using it only when you feel necessary. Do not be frightened to try and use your arm at waist level for light tasks. You will be seeing a physiotherapist and doing regular exercises at home to get the joint moving and to start regaining muscle control. If you feel unsure about what you can or cannot do, please discuss this with the physiotherapist. Lifting your arm in front of you **may** still be difficult at this stage.

Phase 3. Between 6 and 12 weeks

The pain should be lessening. The exercises are now designed to improve the movement available and get the muscles to work, taking your arm up in the air or away from your body when you are sitting or standing. Overall, you will have an increasing ability to use your arm for daily tasks (see driving, work and leisure sections later).

Phase 4. After 12 weeks

You can progress to more vigorous stretches if this is necessary for the activities that you want to do. If the muscles are weak because before the operation the shoulder pain stopped you being able to use them, you should find that you will regain the strength in them with regular exercise. Strength can continue to improve for many months, even up to a year or more. However, unfortunately sometimes the muscles are badly damaged and then you may find it is difficult to regain movement, even though you are trying very hard. Even if the muscles will not work properly, the pain in the shoulder joint should still be much less than before your operation and often you can find small 'trick' movements that enable you to do what you want to do.

Most improvement will be felt in the first 6 months, but strength and movement can continue to improve for 18 months to 2 years.

When can I return to work?

You will probably be off work approximately 6-8 weeks, depending on the type of job you have. If you are involved in lifting, overhead activities or manual work you are advised not to do these for 3-6 months. Please discuss any queries with the therapists or hospital doctor.

When can I drive?

It is normally about 6-8 weeks before you can do this safely. You may find it is more difficult if your left arm has been operated on because of using the gear stick/handbrake. Check you can manage all the controls and it is advisable to start with short journeys. The seat belt may be uncomfortable initially, but your shoulder will not be harmed by it.

In addition, check your insurance policy. You may need to inform the insurance company of your operation.

When can I participate in my leisure activities?

Your ability to start these activities will be dependent on pain, range of movement and the strength that you have in your shoulder following the operation. Please discuss activities in which you may be interested with the therapists or hospital doctor. Start with short sessions, involving little effort and gradually increase.

General examples:

Swimming – after 6 weeks

Gardening (light tasks, eg, weeding) – after 6-8 weeks
(heavier tasks, eg, digging, mowing lawn) after 3-6 months

Bowls – after 3-6 months

Golf, tennis, badminton or squash – after 4-6 months.