

Dressings

On the day after your operation, please remove the outer dressings until you come to a small dressing at the level of the elbow. Please leave this dressing alone and do not allow it to get wet.

If you feel your dressing needs changing before your stitches are removed, please contact your physiotherapist or GP's surgery.

Stitches

If your stitches need to be removed, you should make an appointment with your GP for 10-14 days after your surgery. You will be advised on the day.

Follow-Up Care

Any follow-up appointments will be given to you before you leave or sent to you by post.

If you require this leaflet in any other format, e.g., large print, please telephone
01935 384256

If you have any questions or require further information or advice, please contact one of the following:

DAY SURGERY UNIT - Mon - Fri

(between 8 am and 7 pm) 01935 384339

PHYSIOTHERAPY DEPARTMENT - Mon - Fri

(between 8:30 am and 5 pm) 01935 384358

ORTHOPAEDIC CLINIC - Mon - Fri

(between 9 am and 5 pm) 01935 384319

After 7 pm, the Senior Sister on Duty

(Clinical Site Manager) 01935 384525

ULNAR NERVE DECOMPRESSION

*Day Surgery Unit
and
Hand Therapy Service*

Please read this sheet carefully and if there is anything you do not understand or if you require further information, speak to your nurse or physiotherapist.

ULNAR NERVE DECOMPRESSION

This operation is performed to release pressure on the ulnar nerve at the elbow.

On the ward

You will be visited by the doctor, who will discuss the operation and give you an opportunity to ask any questions.

The site of the operation will be marked and your written consent for the operation obtained. All jewellery on the affected side should be removed.

AFTER THE OPERATION

Discomfort

You will be offered mild pain relief medication to take for the first 2 to 4 days to deal with any discomfort you may have. Severe, continuous pain should be reported to your doctor.

Swelling

To aid the removal of swelling in your hand, it is essential that you follow the instructions below:

1. You will be supplied with a sling or collar and cuff to keep your hand above your heart when standing and walking. Use this for the first 48 hours after surgery.
2. Remove your sling hourly so you can do the exercises described in the next section.
3. When sitting or lying, support your arm and hand on pillows at heart height.
4. If you experience excessive swelling of the hand and fingers, it may indicate that the bandage is too tight particularly if, in addition, the fingers are very painful, numb, cold or blue. If these symptoms are not improved by exercising the fingers as described in the next section, contact the Casualty Department of your local Hospital or your own doctor.

Exercises

It is extremely important that you practice the following exercises for approximately 5 minutes every hour during the day in order maximum movement to be regained following the operation. Slow, controlled exercises are more effective and more comfortable than quick movements. The exercises should be started as soon after the operation as possible:

1. Stretch your arm above your head
2. Bend and straighten your elbow with your arm supported on a table
3. Turn the palm of your hand to face the ceiling and then the floor
4. Bend your wrist backwards and forwards
5. Stretch your fingers and thumb apart as far as possible
6. Bend your fingers to touch the palm of your hand as to make a fist
7. Try to touch the tip of each finger with your thumb